

Volume 1, Issue 2

Research Article

Date of Submission: 30 June, 2025

Date of Acceptance: 12 August, 2025

Date of Publication: 20 August, 2025

A Concept of Energy, Embracing Earth's Environmental Mechanisms

Robert C. Kempton*

Chartered Civil Engineer and Environmentalist, LL.M., Master of Laws, Environmental Law and Practice, United Kingdom

***Corresponding Author:** Robert C. Kempton. Chartered Civil Engineer and Environmentalist, LL.M., Master of Laws, Environmental Law and Practice, United Kingdom.

Citation: Kempton, R. C. (2025). A Concept of Energy, Embracing Earth's Environmental Mechanisms. *J Rehabil Res Curr Updates*, 1(2), 01-12.

Abstract

The universal force of attraction between all matter dominates Earth's 'environmental mechanisms' through the Moon's orbit and Man's concept of gravitational forces, creating intricate relationships between energy flow and the planet's ecosystems, such as photosynthesis, geothermal activity, biogeochemical movements and hydrological cycles. These dynamic reactions shape both the stability and evolution of the planet's systems. The Moon's gravitational pull regulates tidal movements, influencing marine ecosystems and coastal landscapes. Similarly, the intricate balances within biogeochemical cycles ensure the seamless transfer of nutrients between living organisms and the inert components of Earth's surface, sustaining life across the biosphere.

The Oceans cover over 70% of Earth's surface and are in perpetual motion due to earth's natural environmental mechanisms, providing vast natural 'energy storage' reservoirs for major kinetic energy resources around the globe. The development of Man's Science 'fluid mechanics' techniques, to harness this energy, will provide long-term sustainability for mankind.

Human activities are now impacting on all major ecosystems and the biogeochemical cycles they influence, and urgently requires careful identification and analysis. This requires new international laws on Climate Change and other related issues causing severe harm to all matter on our planet.

Man's present approach to nature's biogeochemical cycles for Earth's 'matter' must be addressed, particularly related to present laws of Physics and Chemistry, and with reference to the modern framework for the Periodic Table.

Keywords: Environmental Mechanisms, Fluid Mechanics, Boundary Conditions, Stratification, Kinematic Viscosity, Energy, biogeochemical process, Matter, Periodic Table

Appendices

[To be read in conjunction with]

Appendix 1: Fluid Mechanics

Appendix 2: Additional Methodologies / Regional High Seas and Oceans

Addendum

[for general reference]

1.) Periodic Table – Basic format / structure

2.) Structure of the Atom

3.) Atmospheric Prevailing Winds

Introduction

The universal force of attraction between all matter dominates Earth's environmental mechanisms, with the Moon's location prominent.

This gravitational concept embraces the intricate relationships between energy flow and the planet's ecosystems, such

as photosynthesis, geothermal activity, biogeochemical cycles, and hydrological cycles. Indeed, it provides an approach towards a better understanding of these mechanisms in a sustainable manner.

Atmosphere

The energy from the Sun heats the atmospheric matter, which rotates in unison with Earth and is responsible for creating the circulation of all 'elastic fluidised' ¹ matter. The curvature of Earth creates the variation of heat between the equator and the poles, producing two atmospheric (Hadley) cells, less dense air at the equator rises creating low pressure, and cold dense air falls at the poles.

The environmental mechanism on earth allows the air to rise at the equator and flow to the 30° latitudes, creating three cells for each hemisphere, (Hadley / Ferrel / Polar)². [Ref: Addendum 3]

The earth's rotation on a tilted axis, produces perpetual motion of the oceans in the southern hemisphere, while there is a greater land mass in the northern hemisphere. This creates complex global weather patterns across the six cells around the planet. The dynamic of combined mechanism switching between temperature and pressure, drives the movement of air masses, influencing weather patterns and climatic zones. The atmospheric mechanisms also facilitate the distribution of moisture, shaping ecosystems and weather systems across the planet, including atmospheric pollution³.

Anthropology defines the Coriolis effect, resulting from earth's rotation, as the pattern of deflection taken by objects not firmly connected to the ground as they travel long distances around earth, and which is responsible for many of the critical weather patterns. At the equator a maximum distance is travelled in one day, approaching 1,000 miles⁴, while the distance travelled at the poles, approaches zero travel. In the northern hemisphere the motion is to the east, while in the southern hemisphere it is to the west. The build-up of trade winds and cyclones are examples of the Coriolis effect.

Oceans⁵

The Sun's energy is directly responsible for the stratification process to earth's oceans, which creates the environmental mechanisms to produce layers of transition boundaries through the ocean's depths, vital for all marine life, as we know and understand it. In Lakes and small bodies of water, the warm surface layer is less dense and forms a thermocline boundary condition, above which is the 'epilimnion', with the cooler water disconnected below as the 'hypolimnion'.

In the Oceans, the thermocline's environmental mechanism is influenced by the depth of the ocean's waves, which creates a mixing process boundary layer condition in the upper 100 metres. The temperature decreases rapidly from the mixed upper layer of the ocean (epipelagic zone) to the much colder deep water in the thermocline (mesopelagic zone).

Below the thermocline are colder, denser waters, and below 3,300 feet to a depth of about 13,100 feet, the water temperature remains constant⁶. The deeper (stratified) layers contain colder, denser bottom waters where rare and unknown life forms exist⁷.

The relationship between depth and temperature is significant, as different marine species thrive in specific layers. This environmental mechanism may be a key area for studying the origins of life.

Creatures who live at great depths do not have air in their bodies such as the swim bladders found in fish that live in more shallow waters. Without air in their bodies, the pressure problem is solved. Fish, crab, octopus, worms, limpets and clams are just some of the creatures found in the depths of the oceans.⁸

Despite its central role in the global climate, the Southern Ocean circulation is still one of the least understood ocean circulation systems of the planet⁹.

The term El Niño (Spanish for 'the Christ Child') refers to a warming of the ocean surface, or above-average sea surface temperatures, in the central and eastern tropical Pacific Ocean¹⁰. The low-level surface winds, which normally blow from east to west along the equator ("easterly winds"), instead weaken or, in some cases, start blowing the other direction (from west to east or "westerly winds"). El Niño recurs irregularly, from two years to a decade, and no two events are exactly alike. El Niño events can disrupt normal weather patterns in the United States and globally.

The structural intricacies of ocean layers are further influenced by nutrient cycles and biological activity. In the epipelagic zone, where sunlight penetrates most effectively, primary producers like phytoplankton perform photosynthesis, generating the foundation of oceanic food webs. These upper layers are teeming with life due to nutrient upwelling, a phenomenon driven by ocean currents and the movement of tectonic plates. As depth increases, the availability of sunlight diminishes, creating stark contrasts in biodiversity and energy dynamics between the epipelagic and mesopelagic zones.

The mesopelagic zone, often referred to as the "twilight zone," hosts unique adaptations among its inhabitants, such as bioluminescence and specialized feeding mechanisms, enabling survival in low-light conditions.

This zone also serves as a transitional environment where carbon and nutrients are sequestered, playing a critical role

in the global carbon cycle.

For example, phytoplankton, which are tiny plant-like organisms, flourish in sunlight-rich zones, whereas, zooplankton, which feast on these nutrients, occupy various depths depending on the time of day.

The 'high seas' is in perpetual motion, embracing vast kinetic energy resources¹¹, sufficient for long-term sustainability, and the Oceans are earth's 'natural Energy Storage reservoirs' for all mankind. Man's Science is now sufficiently developed to co-ordinate and organise a Structure Plan embracing the additional methodologies¹² to harness all earth's natural energy resources, based on 'Fluid Mechanics' concepts.

New Environmental Laws¹³ to embrace the high seas vast energy resources are beyond the scope of this paper, but mention should be made that the high seas remain beyond national jurisdiction, at this time, and the new laws, when ratified, will require expansive amendments in future to embrace these natural energy resources.¹⁴

Climate change¹⁵

is a part of 'Earth's environmental balance mechanism', defined by 'boundary conditions' and based on 'Man's Laws of Science'? Human activities are now impacting all major ecosystems and the biogeochemical systems they influence, necessitating careful identification and analysis. 'Earth Systems Science', encompassing geology and chemistry, pertains in part to 'the movement and transformation of chemical elements and compounds between living organisms', known as the biogeochemical cycle, which involves 'fluxian'¹⁶ boundary conditions through anthropological boundaries, and land matter conditions.

Environmental Mechanisms, embracing Greenhouse Gases (GHG).

Energy enters or leaves the Earth's environmental system through electromagnetic radiations. It radiates from the surface by 'infrared radiation', 'visible light' and 'ultraviolet radiation'. This temperature is controlled by the planet's environmental mechanisms and when Earth's surface absorbs as much energy as it releases, the planet's energy budgets are balanced, and the temperature remains stable.¹⁷ The following points apply:

- Extremely high energy objects (Sun) emit high-energy electromagnetic radiation, classified as short wavelength, embracing 'ultraviolet radiation' and visible light.
- For cooler energy objects, such as Earth's surface, much less intense radiation has a longer wavelength radiation and is classified as infrared radiation.
- Earth's atmosphere is not totally transparent to all electromagnetic radiations, it behaves as a selective filter, which absorbs, reflects and allows through distinctive radiations bands, dependent on the atmospheric make-up of the gases.
- The physical properties of some gases tend to be opaque to infrared radiations. When present in the atmosphere, they reflect a proportion of gases emitted from earth back towards the surface, and this causes an imbalance, warming the earth's surface, both day and night. It prevents some of the heat absorbed from the Sun from being released back to space.
- This balancing environmental mechanism appears to be unique to our small planet, Earth. The mechanism is critical to life. Without the 'partial opacity' of Earth's atmosphere to infrared radiation, the planet's surface would reach sub-zero temperatures at night when cut-off from sunlight. Life forms would cease, as on the Moon.

Carbon dioxide is one of the primary contributors to the 'greenhouse effect', which traps heat within Earth's atmosphere. Its sources are varied, ranging from natural processes such as volcanic eruptions and respiration to human activities including the burning of fossil fuels and deforestation. The environmental impact of CO₂ extends beyond the temperature rise condition it influences ocean acidification, disrupts ecosystems, and accelerates biodiversity loss.

Matter is anything that occupies space, has mass, and is made up of substances called elements, which are composed of extremely small particles called atoms, and have specific chemical and physical properties. Man's Science classifies three 'body states' of matter as, gaseous / fluid / solid states, with the Science 'Fluid Mechanics'¹⁹, embracing gaseous as elastic fluids.

Man's Science explains the atom structure as comprising, protons, neutrons, and electrons, perceived within a sphere of influence, controlled by 'attraction' and 'repulsion' forces, and not by a boundary condition around that perceived sphere of influence.

Size (Space): Man's Science has no 'real perception' of size related to the world's surroundings and our environment. He is unable to conceive the sheer magnitude of the Universe when related to the conceived size of the perceived atom, and is beyond the scope of this paper, except to mention a general observation in Addendum, 2.

The complete tabular array of all chemical elements is classified under the 'Periodic Table'²⁰. The structure co-ordination of the chemical elements, organized by atomic number, is from the element with the lowest atomic number, hydrogen, to the element with the present highest atomic number, 'oganesson' (Fig 1). Reference Addendum 1- Important Basic

Facts of present Periodic Table.

The fundamental elements interact dynamically to form molecules, which are the building blocks of matter. For example, water (H₂O), a critical molecule for life, exhibits unique properties like high 'specific heat' capacity, 'cohesion', and 'surface tension', which are essential for regulating Earth's climate and supporting ecosystems.

Fluid Mechanics²¹

Under man's perceived mechanism, if a fluid is moving slowly, within a constrained space, a shear stress is created within the liquid called 'viscosity'. Sir Isaac Newton²² referred to this as the 'coefficient of molecular viscosity'²³.

"All forms of life on earth are immersed in a fluid or another, either the air of the atmosphere or the water of a river, lake or ocean; even, soils are permeated with moisture. So, it is no exaggeration to say that life, including our own, is bathed in fluids. A slightly closer look at the situation further reveals that it is the mobility of fluids that makes them so useful to the maintenance of life, both internally and externally to living organisms".²⁴

The study of 'matter' and its transformations²⁵ is not merely academic; Man's Laws of Science needs to underpin technological advancements and solutions to pressing global challenges, such as renewable energy and long-term sustainable resource management.

Energy / Matter Interactions:

Energy, the capacity to do work, is intricately connected to matter. In physics, the 'Law of Conservation' states that energy cannot be created or destroyed but only transformed. This principle governs natural processes and Classifications²⁶, from the photosynthesis that powers ecosystems, through biogeochemical processes, and to the geothermal activity driving tectonic shifts.

Fresh approaches towards a long-term Sustainable future: The connection between energy and matter is crucial in understanding Earth's systems and environmental mechanisms, as it determines the flow of nutrients, the cycling of carbon, and the dynamics of climate regulation.

Early History of the Elements and Original Periodic Table

Mendeleev initiated the famous periodic law that Element properties are a periodic function and to organize the elements according to their atomic weight (later changed to Atomic No), while also leaving gaps for undiscovered elements that he predicted based on the periodicity of the table. This prediction was later confirmed with the discovery of gallium, scandium, and germanium, which had properties that matched Mendeleev's predictions.

Johann Wolfgang Dobereiner – The law of Triads 1829 - He observed that groups of three elements (triads) could be formed in which all the elements shared similar physical and chemical properties.

John Newlands, British, 1864 – first to arrange the elements into a periodic table with increasing order of atomic masses. He Identified pattern repeated every 8th element – 'law of Octaves'. Mendeleev (1834–1907) – Was considered the first for preparing a Framework Plan to set-up and record the idea of co-ordinating a structured range of elements (1869).

He wrote Books on the principles for Chemistry, including:

- Organic Chemistry (carbon compounds & living things)
- Inorganic Chemistry (minerals.)

Julius Lothar Meyer (1830–1895) – He arrived at the same theory as Mendeleev. They lived over 1000 miles apart, in the same period. This synchronicity in elements discoveries raises questions about how shared knowledge and scientific exchanges of the time, like those at Heidelberg, could foster parallel development.

Ernest Rutherford (born August 30, 1871, was the first person to split the atom (1918). He is known for his pioneering studies of radioactivity and the atom. He discovered that there are two types of radiation, alpha and beta particles, coming from uranium.

Early History of the Elements and Original Periodic Table:

Mendeleev initiated the famous periodic law that **Element properties are a periodic function** and to organize the elements according to their atomic weight (later changed to Atomic No), while also leaving gaps for undiscovered elements that he predicted based on the periodicity of the table. This prediction was later confirmed with the discovery of gallium, scandium, and germanium, which had properties that matched Mendeleev's predictions.

Johann Wolfgang Dobereiner – The law of Triads 1829 - He observed that groups of three elements (triads) could be formed in which all the elements shared similar physical and chemical properties.

John Newlands, British, 1864 – first to arrange the elements into a periodic table with increasing order of atomic masses. He Identified pattern repeated every 8th element – 'law of Octaves'.

Mendeleev²⁸ (1834–1907) – Was considered the first for preparing a Framework Plan to set-up and record the idea of co-ordinating a structured range of elements (**1869**). He wrote Books on the principles for Chemistry, including:
- Organic Chemistry (carbon compounds & living things)
- Inorganic Chemistry (minerals.)

Julius Lothar Meyer²⁹ (1830–1895) – He arrived at the same theory as Mendeleev. They lived over 1000 miles apart, in the same period. *This synchronicity in elements discoveries raises questions about how shared knowledge and scientific exchanges of the time, like those at Heidelberg, could foster parallel development.*

Ernest Rutherford³⁰ (born August 30, 1871, was the first person to split the atom (1918). He is known for his pioneering studies of radioactivity and the atom. He discovered that there are two types of radiation, alpha and beta particles, coming from uranium.

Figure 1

Global Long-term Sustainability Project (embracing environmental mechanisms)

At last, Man has the knowledge to develop new innovative methodologies required to harness earth's natural energy resources. Vast kinetic energy reserves are available in the high seas, which cover over 70% of the surface of earth.

An early example of Man's Innovative ideas, was recorded in the late 60s, following research work into 'fluid mechanics' relating to 'thick liquids' for municipal sewage sludges. Research studies²⁷ were successfully carried out on site to ascertain the effective range of 'kinematic viscosity' of 'thick liquid' municipal sludges, embracing the importance of 'boundary conditions', and 'shear stress' mechanisms within a fluid. (Table 1) The kinematic viscosity of water was recorded at 0.000012 ft²/sec. (Reference Appendix 1) A research paper²⁸ was presented to the UK, Institution of Civil Engineers in 1971 and awarded recognition in July 1972. The results were adopted in the Leicester City Corporation's Sludge Disposal Project design works, to replace temporary facilities with a permanent structure, adjacent to their Compost Plant, and was successfully completed in 1970.

Table 1 furnishes the range of 'Kinematic Viscosity' figures used in the Design Works calculations for their successful "Sludge Tankering Project" at the Wanlip Sewage Treatment Works.

% solids	Calculated Kinematic Viscosity	Required pipe size	Time for 1,000 galls to flow to reservoir
2.39	.0007 ft ² /sec	Transitional	Zone
2.67	.001 ft ² /sec	-	-
4.27	.00552 ft ² /sec	6" dia	22.85 mins.
4.27	.00552 ft ² /sec	8" dia	7.12 mins.
4.27	.00552 ft ² /sec	12" dia	1.44 mins.
4.27	.00552 ft ² /sec	15" dia	.585 mins.
4.45	.011 ft ² /sec	6" dia	46.00 mins.
4.45	.011 ft ² /sec	8" dia	14.32 mins.
4.45	.011 ft ² /sec	12" dia	2.88 mins.
4.45	.011 ft ² /sec	15" dia	1.175 mins.

Range of Kinematic Viscosity values for Municipal Sludge
 - Wanlip STW Sludge Project, 1971
 (Imperial Units)

Table 1

With the Advent of the 'Biodiversity Before National Jurisdiction' (BBNJ)²⁹ Agreement being ratified, further important legislation is now required to embrace the additional methodologies for harnessing the natural kinetic energy storage in the Regional High Seas.

This Agreement recalled the relevant provisions of the United Nations Convention on the Law of the Sea (UNCLOS)³⁰ of 10 September 1982 and covered just the global surface for the territorial sea and its contiguous zone³¹.

New legislation must include the regional areas of the High Seas for coastal nations, and to be embraced under the UNCLOS 'high seas sections'³², to embrace the methodologies for harnessing natural kinetic energy storage zones.

One methodology to identify is in the 'Epipelagic Zone'³³ of the high seas, which could embrace the traversing of sailing vessels over vast areas. Fluid mechanics design techniques to be adopted to provide hulls of ships to be fitted with equipment during their passage journeys through the surface waters, to 'collect / Hoover / transfer / store' the continuous kinetic energy produced. Major Energy Companies³⁴ to be set up to prepare schedules for multiple passages across the regional seas, with destinations to many energy ports within the coastal management zones.

Slope Currents contiguous to the Continental Shelves.

- Japan's Kuroshio currents: Japan's coastline, due to unstable Teutonic plates, appear unsuitable for fixed turbines to the seabed. Reference 'Experimental verification of a floating ocean current turbine with single rotor for use in the Kuroshio current.'³⁵

- European "Slope Current around the British Isles, potential location for European Slope Current Project, embracing stabilised bathymetric seabed profile³⁶.

Summary, and linkages to Relevant Environmental References

The Incompleteness and uncertainties of Man's Laws of Science towards a Long-term Sustainable Future have revealed much about the complexities of Earth's systems, and they also underscore the fragility of our small planet. By leveraging the principles of matter, energy, and environmental mechanisms, a sustainable future may be envisioned. This involves not only technological innovation but also a harmonious integration of human activities with the planet's natural cycles, ensuring that the balance essential for life is maintained for generations to come.

Pressure: Relentless increase in 'pressure's mechanism' throughout Earth's Ocean depths produce multiple linkage to other mechanisms, creating ever evolving environmental conditions which may be key areas for studying the origins of life as we currently understand it. Life's stratified layers, down to the Mariana Trench depth, are vital research areas for mankind.

It is now essential to restructure an intelligent approach towards setting out a UK structured framework for 'air pollution' controls based upon 'common but differential responsibility'³⁷.

Vast Kinetic Energy resources are available, once additional methodologies have been developed, to harness earth's natural energy resources, as it will provide a long-term sustainability for all mankind, once the World's legislation has been

ratified and brought into force³⁸. An approach towards a Structure Plan Model for Oceans total energy methodologies is recommended, based on proven modern project management services techniques. (APMS)³⁹.

Fluid Mechanics embraces the major 'environmental mechanisms'⁴⁰ within 'Man's Science'.

The protocol on environmental protection to the Antarctic treaty to be considered as a suitable model⁴¹.

Definition

"A fluid is matter in a readily distortable form, so that the smallest unbalanced external force on it causes an infinite change of shape, if applied for a long enough time".

Newton's Laws of Motion

1st Law: Motion of object remains constant unless a force act upon it.

2nd Law: force of an object is equal to its mass times its acceleration

3rd Law: two objects apply forces to each other, equal magnitude / opposite direction.

A Newtonian Fluid, such as water is perceived to operate in the 'turbulent flow' zone range, with sufficient velocity to create the turbulent state. Reference: 'Thin Liquids'.

Materials with different physical properties, are termed Non-Newtonian Fluids. Dense 'thicker' liquids are perceived under 'laminar flow' range. Reference: 'Thick Liquids'. Viscosity: fluid's resistance to flow, (thickness of a fluid.)

Viscosity calculated in the Laboratory, known as Absolute Viscosity

Dynamic density calculated in the field, called the kinematic viscosity.

Boundary Layer conditions create Inertia Force on fluid in motion.

Reynold's No = Ratio of Forces perceived in fluid motion, [Inertia force / viscous force.]

Note: Can only be used for one type of boundary face, such a rugosity of circular pipe, or hull of ship motion in water, or shape of plane wing in flight. Only perceives rugosity to particular shape of boundaries.

'Boundary Layer': Influence on fluids flowing in circular Pipes:

'Thick Liquids' – Laminar flow zone properties – Re No. below 2300 Reynolds Number (applicable to flow in pipes only).

Smooth flow conditions through circular pipes. C/S pipe velocities: 0 to Maxm in Centre. Flow conditions perceived as shear stresses on very large number of the 'shear rings' motion through its contiguous shear rings.

'Thin Liquids' – Turbulent flow zone properties set of curves based on bore roughness

Universal pipe friction diagram chart, adopted for pipes of all diameters and roughnesses, and for all velocities and 'kinematic viscosities. The uncertain theory requires a transitional zone chequered pattern, between the 2 zones. See Diagram 1.

Laminar Flow: Poiseuille law or Poiseuille equation, is a physical law that gives the pressure drop in an incompressible and Newtonian fluid in laminar flow flowing through a long cylindrical pipe of constant cross section.

Turbulent Flow: Darcy Equation is a theoretical equation that predicts the frictional energy loss in a pipe based on the velocity of the fluid and the resistance due to friction. It is used almost exclusively to calculate head loss due to friction in turbulent flow.

End Note

Temperature has a major effect on the viscosity of any fluid.

Information Sheet

26 June 2025

Subject: Additional Methodologies to harness Ocean Energy in the Regional High Seas⁴²

Augmentation of present Methodologies to harness vast kinetic energy resources in the High Seas⁴³.

Proven performance of existing methodologies (Continental Shelf)

(1. La Rance⁴⁴ Tidal Power Station / 2. MayGen⁴⁵ Tidal Power Station)

- Tidal Range Projects in coastal waters

Sihwa Lake Tidal Power Station is the world's largest tidal power installation, with a total power output capacity of 254 MW Megawatt. When completed in 2011, its surpassed France's 240 MW Rance Tidal Power Station, which was the world's largest for 45 years. It is operated by the Koreann Water Resources Corporation.

- Tidal Stream Projects in coastal waters

MayGen Tidal Energy Project, continues to break records, delivering over 37GWh of clean and predictable renewable energy⁴⁶.

"In December 2024, the final turbine of the four turbines for Phase 1 were deployed, and therefore the site is now fully operational. This means that the site is delivering 6 MW of power"⁴⁷.

Classification for High Seas Expected Methodologies to Harness Kinetic Energy Resources in Oceans.

Epipelagic Zone (mixed layer): Ocean Passage collection projects in regional high seas.

Reference 6.2. (i)

Mesopelagic Zone – restricted to transition zone for carbon sequestration.

Slope currents contiguous to continental shelves.⁴⁸

Continental Slope Stream Projects

Arctic Ocean⁴⁹ Deepwater current along Oceans' transition zone

Deep Stream Oceans Projects

References

Energy Act 2023 /footnote, Addendum A, p24

Reference, Part I Knew Domestic Law, Patentability, Inventive step

118 Atoms are Classified in the form of 7 Periods (Rows) and 18 Groups (Columns)

To follow the perceived structure of physical statistics, two strings named lanthanides and actinides are located below in 2 rows

The lanthanide or lanthanoid series of chemical elements comprises at least the 14 metallic chemical elements with atomic numbers 57–70, (element 71), is sometimes included⁵⁰

The actinide or actinoid series encompasses at least the 14 metallic chemical elements also included with atomic numbers from 89 to 102, lawrencium is sometimes included.

Distinctive Groups

The Alkali Metals (Group 1, except hydrogen)

The Alkaline Earth Metals (Group 2)

The Transition Metals (Groups 3-11)

Post Transition Metals

located between transition metals and metalloids.

Metalloids

display both properties of metals and non-metals.

Halogens Group 17

Fluorine, F Chlorine, Cl Bromine, Br Iodine, I & Radioactive elements Astatine, At Tennessine, Ts

Noble Gases Group 18

Helium, He neon, Ne argon, Ar krypton, Kr xenon, Xe radon, Ra oranesson, Og---- Group 18 (VIIIa) ---- Initially believed they could not combine with other elements. (Later, Group No. changed from group O to group 18.)

Basic Structure Spine Framework

The outline structure frame is perceived as being presented in 4 blocks – s-d-p-f.

The f-block, 'fundamental' intended before column 3, and the two rows shown below blocks s, d, p.⁵¹ (Lanthanide and actinide rows.)

End notes

Metals are found on the left of the periodic table and non-metals on the right.

Man's Science perception for Chemical elements structure plan⁵² (Reference Addendum 1) The periodic table Structure may be used to locate any element.

Anthropology States the Elements of life as

carbon (C), oxygen (O), hydrogen (H), and nitrogen (N)—which make up 96% of living matter.

Main Components of air N, 78%, O₂ 21%, CO₂ 0.04%, Argon .93%, Traces .03%

Perceived Definition

Atoms consist of a nucleus containing protons and neutrons, surrounded by electrons in shells. The numbers of subatomic particles in an atom can be calculated from its atomic number and mass number⁵³.

Man's Science

Atoms are perceived to be extremely small, positively charged nucleus surrounded by a cloud of negatively charged electrons. The nucleus is perceived to be less than one ten-thousandth the size of the atom the nucleus contains more than 99.9% of the mass of the atom⁵⁴.

Size

A convenient unit of length for measuring atomic sizes is the angstrom (Å), defined as 10^{-10} metre. The radius of an atom measures 1–2 Å. Compared with the overall size of the atom, the nucleus is even more minute. It is in the same proportion to the atom as a marble is to a football field. In volume the nucleus takes up only 10^{-14} metres of the space in the atom—i.e., 1 part in 100,000. A convenient unit of length for measuring nuclear sizes is the femtometre (fm), which equals 10^{-15} metre. The diameter of a nucleus depends on the number of particles it contains and ranges from about 4 fm for a light nucleus such as carbon to 15 fm for a heavy nucleus such as lead. Despite the small size of the nucleus, virtually all the mass of the atom is concentrated there.

Perceived Structure: The protons and neutrons form the atom's central nucleus.
(The ordinary hydrogen atom is an exception; it contains one proton, but no neutrons.)
Protons have a positive electrical charge.

Neutrons are electrically neutral—they carry no charge.
Circling the nucleus is a cloud of Electrons, which are negatively charged.
Perceived like opposite ends of a magnet that attract one another, which binds them to the nucleus.

The nucleus is small and dense compared with the electrons, which are the lightest charged particles in nature.

The electrons circle the nucleus in orbital paths called shells, each of which holds only a certain number of electrons. The outer shell is the valency shell.

An ordinary, neutral atom has an equal number of protons (in the nucleus) and electrons (surrounding the nucleus); thus, the positive and negative charges are balanced.

Some atoms, however, lose or gain electrons in chemical reactions or in collisions with other particles.

Ordinary atoms that either gain or lose electrons are called ions. If a neutral atom loses an electron, it becomes a positive ion. If it gains an electron, it becomes a negative ion.

These basic subatomic particles—protons, neutrons, and electrons—are themselves made up of smaller substances, such as quarks and leptons.

As Perceived through quantum physics: The electron wave is uniform in all directions from the nucleus, is peaked at the centre of the atom, and has the same phase everywhere. Higher energy levels in the atom have waves that are peaked at greater distances from the nucleus.

Quantum Physics definition: Atoms, the smallest particles of an element that exhibit the properties of that element, consist of negatively charged electrons around a central nucleus composed of more massive positively charged protons and electrically neutral neutrons.^{23 Jun 2019}

Endnote

Perceived as a shell, or wave, or created, or formed by another undiscovered energy form?

Simplest Model




Figure 1




Figure 2




Figure 3

Table of Legislation

Climate Change Act 2008, UK Public General Acts, 2008 c.27

UNCLOS - Agreement under the United Nations Convention on the Law of the Sea on the Conservation and Sustainable use of Marine Biological Diversity of Areas Beyond National Jurisdiction, New York September 2023 [The Agreement is not yet in Force]

UNCLOS, Oceans and Law of the Sea, Division for Ocean Affairs and the Law of the Sea, United Nations Convention on the Law of the Sea of 10 December 1982

Patents Act 1977, UK Public General Acts, 1977c. 37, Part 1, Patentability, 4 Industrial application. Energy Act 2023, UK Public General Acts, 2023 c. 52.

References

Bell S et al, Environmental Law, Ninth Edition, 2017, Air and atmospheric pollution, p 530-538 Cushion-Roisin B et al, 1st Edition, 2008, "Fluid Mechanics of Environmental Interfaces", ABSTRACT – Chapter, Environmental fluid mechanics: Current Issues and future outlook.

Cushman-Roisin B et al "Environmental Fluid Mechanics: Current issues and future outlook", 1st Edition, 2008, Abstract, 1st paragraph Darcy, 17th century Paris: Darcy measured the discharge of a tube by measuring the volume of water that flows out of the outlet during a fixed period of time.

Earth System Science, Earth Labs, Lab 4: It's all connected; Global Circulation, Part A: Tracing Pathways, Caption Energy and Power Supply Statistics, Forms of known Energy / Addendum A, Information Sheet, - Energy / Addendum B

-Information Sheet – Energy storage.

Flow Conditions in Pipes due to Thick Liquids, 1971. Paper presented at the North-West Institution of Civil Engineers Student and Graduate Meetings in March 1971, ICE Miller Prize Award recognition in July 1972. – Open Research work, 1971, referenced in this paper

Fluid Mechanics definition and terminology, Appendix 1:

Fluid Mechanics: At that time literature and knowledge available was somewhat limited to movement of 'thin liquids' through circular pipes. Flow charts were readily available to the student or engineer, giving a variety of curves for flow through pipes of various roughness, age, and material.

Francis J.R.D, A Textbook on Fluid Mechanics, The Properties of Fluids, 2nd Edition 1962, 1.1 Definition of a Fluid.

Francis J.R.D, A Textbook on Fluid Mechanics, The Properties of Fluids, p 2-3 / The pipe friction diagram for pipes, Fig. 13.10, p207

Francis J.R.D, B Sc (Engineering) London, Assoc. I.C.E, with Contribution on Gas Flow from Jackson M.A. (Cantab), A Textbook of Fluid Mechanics", 1958, p3, top of page.

Geoinformatics & Geostatistics, A Scientific Journal, SciTechnol: An Overview, Review Article, 2025, 13.4, p6

Glossary of Archaic Chemical Terms, Part II (C-F), elastic fluid definition.

Gwam CU (2010) World Meteorological Organization (WMO), Technical references.

Howarth W, McGillivray D, Water Pollution and Water Quality Law, 2001, Chapter 18 International Marine Environmental Law.

Hughes D et al, Environmental Law, fourth edition, Reference to Chapter 17, Atmospheric pollution, pages 543-578

International Journal Evolving Sustainable and Renewable Energy Solutions, Prime Open Access publication, Vol 1, Issue 1, 20 March 2025, The High Seas in Perpetual Motion, para 1, p7,

Japan's Kuroshio currents (hindered by unstable Teutonic plate conditions

Kempton, Robert, Approach Towards a Total UK Environmental Energy Master Plan for Clean Renewable Energy, Reviewing the Incompleteness of the British Energy Security Strategy after Brexit, and Embracing Tidal Waters and other Innovative Sources (May 17, 2024). Available at SSRN: https://papers.ssrn.com/sol3/papers.cfm?abstract_id=4831998

La Rance Tidal Power Station – Appendix 2. M Auger, P Prandi, JB Sallée - Scientific Data, 2022 - search.ebscohost.com

Mayer Benoit, The International Law on Climate Change, Cambridge University Press, 2018, Introduction, p 2, 1st para.

MayGen Tidal Power Project – Information Sheet – Appendix 2.

NASA, 'Glen Research Center' 27 June 2024 - Sir Isaac Newton worked in many areas of mathematics and physics. He developed the theories of gravitation in 1666 when he was only 23 years old. In 1686, he presented his three laws of motion in the "Principia Mathematica Naturalis."

National Geographic, Education, The Coriolis Effect: Earth's Rotation and its Effect on Weather.

National Weather Service (.gov) <https://www.weather.gov> > winds > Flight Environment, Prevailing Winds.

Newton Issac, Sir, Expression used to explain 'particular boundary' condition. / Reference also Glossary of Archaic Chemical Mayer Benoit, The International Law on Climate Change, Cambridge University Press, 2018, Introduction, p 2, 1st para.

Periodic Table - Addendum 1 – Basic Format / Structure

Periodic Table Reference comment: after La(Z=57) filling of 4f-orbital starts which is complete at atomic number 71. They have different properties and are placed separately at the bottom of the structure in f-block.

Poiseuille was a physician who had been trained in physics and mathematics. In 1846, Jean Louis Poiseuille published a paper on the experimental research of the motion of liquids in small diameter tubes.

Reference to Addendum A, Energy Paper titled: "Approach Towards a Total UK Environmental Energy Master Plan for Clean Renewable Energy, Reviewing the Incompleteness of the British Energy Security Strategy after Brexit, and embracing Tidal Waters and other Innovative Sources"

Reference to early Fluid Mechanics Text-Book, "A Textbook of Fluid Mechanics", Francis J.R.D, B Sc (Engineering) London, Assoc. I.C.E, with Contribution on Gas Flow from Jackson M.A. (Cantab), 1958.

Sands P et al, Principles of International Environmental Law, fourth Edition, Chapter 11, Oceans, Seas and Marine Living Resources.

Shen HH et al 2002, books, google.com, "Environmental fluid mechanics: theories and applications"
Structure of the Atom, Addendum2: [Man's Laws of Science conception]
The Role of Oceanic Thermoclines in Marine Biology / <https://www.nature.com> > articles
World Journal of Forest Research, ISSN: 2994-5569, Research Article: 'Approach to Climate Change Sustainability Project, for 'Our Common Future', 25 April 2025, 3.1.1 The Earth's Wind Pattern Forces Volume 4, Issue 1, p5.

References

1. <https://www.bbc.co.uk/bitesize/guides/zwn8b82/revision/3>
2. <https://www.britannica.com> –
3. <https://www.britannica.com/biography/Ernest-Rutherford>
4. <https://www.britannica.com/science/atom>,
5. <https://origins.osu.edu/milestones/mendeleev-periodic-table-UN-chemistry-radioactivity-noble-gases>
6. <https://www.oxfordreference.com/display/10.1093/oi/authority.20110810105408716>
7. <https://www.bridgew.edu/>
8. <https://www.pmel.noaa.gov/eoi/nemo1998/education/pressure.html>
9. <https://www.chemistrytalk.org>
10. <https://doi.org/10.20944/preprints202312.0085.v1>
11. https://www.weather.gov/source/zhu/ZHU_Training_Page/winds/Wx_Terms/Flight_Environment.htm