Journal of Medical Science and Health Care Research



Volume 2, Issue 1

Research Article

Date of Submission: 12 April, 2025 Date of Acceptance: 02 May, 2025 Date of Publication: 28 May, 2025

Main Clinical Characteristics of Medication Overuse Headache

Srdjan Ljubisavljevic^{1*}, Stefan Todorovic² and Radomir Damjanovic²

¹Faculty of Medicine University of Niš, 18000 Niš, Serbia

²Clinic of Neurology, University Clinical Center Niš, 18000 Niš, Serbia

*Corresponding Author:

Srdjan Ljubisavljevic, Faculty of Medicine University of Niš, 18000 Niš, Serbia.

Citation: Ljubisavljevic, S., Todorovic, S., Damjanovic, R. (2025). Main Clinical Characteristics of Medication Overuse Headache. *J Med Sci Health Care Res, 2*(1), 01-06.

Abstract

Introduction: Medication overuse headache (MOH) is a secondary headache that occurs 15 or more days per month in patients with preexisting headache. This can result from the excessive (10 or 15 days/month) use of drugs for acute or symptomatic headache therapy.

Aim: To examine the relationships between demographic parameters, illnesses and habits of life, clinical type and length of previous headache and MOH and clinical features of MOH.

Materials and methods: This study was conducted at the Clinic of Neurology, Clinical Center of Niš, and included 83 patients (11 men and 72 women) who were first diagnosed with MOH. The mean age of the patients in the study cohort was 40.5 ± 11.6 years. The study was performed in strict accordance with the Declaration of Helsinki after informed consent was obtained from each participant in the study. The study was approved by the local ethics committee.

Results: In the study cohort, there were more women (86.7%), nonsmokers (59%), those who did not consume alcohol (95.2%), those who were physically inactive (90.4%), those who consumed caffeinated beverages (89.2%), and those with diseases of the cardiovascular (12%) or osseous joint system (10.8%). MOH was most commonly generated by chronic migraine (64%). The duration of MOH until diagnosis was 5.1 ± 5.5 years. The following MOH characteristics were observed: moderate pain (45.8%), bilateral localization (56.6%), temporal presentation (65.1%), and neck tightness (81.9%). Common (41%) and combination analgesics (48.2%) were used most frequently, 15 to 25 days per month. The impact of MOH on daily life was assessed as significant (HIT - 65.4 ± 5.5), correlating with the duration of earlier chronic headaches (r = 0.327, p = 0.003).

Conclusion: MOH can be generated by the transformation of a previous chronic headache due to overuse of analgesic therapy. The MOH significantly affects all aspects of a patient's life. MOH can be prevented by educating patients by limiting the use of analgesic therapy, especially in the at-risk group (middle-aged women with chronic migraines).

Keywords: Medication-Overuse Headache

Introduction

Medication overuse headache (MOH) is a secondary headache classified according to the International Headache Society as a group of headaches attributed to the use or discontinuation of various substances [1]. MOH occurs 15 or more days per month in patients with preexisting headaches. It occurs due to regular (at least three consecutive months) overuse (10 days or 15 days for certain medications) of medicines used for acute or symptomatic headache therapy. MOH can occur due to the excessive use of ergotamine, triptans, ordinary analgesics (pracetamol, acetylsalicylic acid, nonsteroidal anti-inflammatory drugs (NSAIDs), opioids, combined analgesics, or simultaneous use of several types of drugs.

This type of headache most often occurs in the fourth decade (between 30 and 50 years of age) and is more common in women (female-to-male ratio is 3- 4:1) [2]. The estimated prevalence of MOH is approximately 3%. The burden on the patient and his family is a consequence of the impairment of the quality of daily life. The MOH represents a significant

burden on the health system and society as a whole and affects the reduction of working capacity, often missing work [1,2].

Although it can occur from any type of headache, MOS is most often caused by the chronic transformation of migraine. The complete mechanism underlying the pathophysiological process that leads to this condition has not been fully elucidated. It is assumed that processes of central sensitization (suppression of endogenous pain modulation, neuronal excitability due to neurotransmitter imbalances [adrenaline, serotonin], etc.) represent the pathogenetic basis of MOH [3-6].

MOH treatment involves educating patients about the necessity of limited use of analgesics, most often with the simultaneous implementation of detoxification therapy and prophylactic support of coanalgetic therapy (valproate, topiramate, amitriptyline, onabotulinum toxin A, etc.). This can be done in an outpatient or hospital setting. During the use of triptans, ergotamine, and NSAIDS, individuals are advised to abruptly stop the excessive use of these drugs. On the other hand, a gradual reduction in the dose of the drug is recommended for previous overuse of barbiturates, benzodiazepines and opioids [7,8].

Although MOH is considered potentially preventable, changeable and unchanging risk factors for MOH have not yet been recognized. Identifying these risk factors could aid in identifying patients with a previous headache who are at higher risk for chronic headache and transforming a previous headache into MOH.

Research Objective

The research aimed to examine the following:

- Associations between demographic parameters, disease status and lifestyle habits, clinical type and duration of previous headache and MOH
- Clinical features of MOH

Materials and Methods

This research was conducted as part of a project approved by the Ethics Committee of the Faculty of Medicine in Niš as an observational, cross-sectional study. The study included all patients examined at the Headache Clinic of the Neurology Clinic, Clinical Center in Nis, during 2019 (January-December), in whom the diagnosis of MOH was made for the first time during this period. Patients gave informed consent to participate in the study. The diagnosis of MOH was made according to the current diagnostic criteria of the International Classification of Headaches [1]. In all patients, the second etiology of the headache was excluded after complete diagnostic processing (computed tomography, magnetic resonance imaging, etc.).

The following data were collected from the included patients: demographic data; a history of personal life; the presence of other diseases; the presence of previous (primary and/or secondary) headaches (type, characteristics, duration of headache, frequency of occurrence, effectiveness of symptomatic and preventive therapy); and habits and risk factors (physical activity, smoking cigarettes, use of alcohol, caffeine, etc.). The data related to MOH included the location of pain (frontal, temporal, parietal, occipital), lateralization (unilateral, diffuse), character (dull, pulsating pain), intensity of the pain (using a numerical scale to assess pain), the existence of associated symptoms and signs (nausea/vomiting, photophobia, phonophobia, diplopia, neck and shoulder tightness: blurred vision, ringing in the ears: hyperacusis), type of analgesic therapy used, preventive therapy, frequency of use of this therapy (number of days/month), effectiveness of the therapy, and effect of headaches on the activities of everyday life (headache impact test - HIT). For the statistical processing of the data, a single database was constructed in which descriptive and analytical statistics were applied. All the statistical analyses were carried out with the SPSS 16.0 software package.

Results

The study included 83 patients with MOH—11 men and 72 women. The mean age of these patients was 40.5±11.6 years (range: 18 to 71 years). By analyzing the frequency of the parameters studied, it was observed that, in the cohort of patients with MOH, there were more women, nonsmokers, and nondrinkers (daily use/use a greater number of days per week); they were physically inactive (daily/greater number of days per week); and they had comorbidities in the form of cardiovascular, bone and joint diseases (Table 1).

Parameter	Number (%) /X±SD			
	Smoking			
Yes		34 (41,0)		
No		49 (59,0)*		
Length of smoking (age)		7,5±11,1		
Number of cigarettes		5,4±7,7		
Alcohol use				
Yes		4 (4,8)		

No	79 (95,2)*			
Caffeine intake				
Yes	74 (89,2)*			
No	9 (10,8)			
Physical activity				
Yes	8 (9,6)			
No	75 (90,4)*			
Comorbidities				
Yes	36 (43,4)			
No	47 (56,6)*			
Comorbidities (Dg)				
Endocrinological diseases	al diseases 7 (8,4)			
Psychiatric Diseases	2 (2,1)			
Neurology Diseases	1 (1,2)			
Cardiovascular diseases	10 (12,0)*			
Respiratory Diseases	7 (8,4)			
Bone and joint diseases	9 (10,8)*			
* Hi square test, p<0.05.				

Table 1: Lifestyle Habits, Risk Factors, and Comorbidities

MOH was more common in migraine patients (64%) than in patients with tension-type headache (25%) or symptomatic headache (11%). The duration of the primary headache was 6.5 ± 3.2 years (Figure 1).





In the study group, the average duration of MOH (up to the time of diagnosis) was more than 5 years, more than half a month was affected by headache, and the duration of individual headache attacks was more than 10 hours per day. Patients described the pain as moderate to severe (45.8%), bilateral (56.6%), mostly temporal (65.1%), or most often accompanied by complaints in the form of neck tightness (81.9%). Most patients used ordinary analgesics (NSAIDs) (41%) or combined analgesics (48.2%), averaging 15 to 25 days per month. This treatment was ineffective in the largest percentage of patients (78.3%). A large number of patients used prophylactic therapy and antidepressant drugs (83.1%). The effect of headaches on activities of daily living was assessed as significant (HIT - 65.4±5.5) (Table 2).

Parameter	Number (%) /X±SD		
Duration (year)		5.1±5.5	
Number of days/months		19.4±6.1	
Duration (h)		10.7±1.9	
Pain (VAS)			
5		16 (19,3)	
6		17 (20,5)	
7		38 (45,8)*	
8		7 (8,4)	
9		3 (3,6)	

10		2 (2,4)
	Localization	
Frontal		14 (16,9)
Occipital		5 (6,0)
Parietal		10 (12,0)
Temporal		54 (65,1)*
	Lateralization	
Bilateral		47 (56,6)*
Unilateral		36(34, 4)
	The character of pain	
Pulsating		41(49, 4)
Dull		42(50, 6)
	Associated symptoms	
photophobia / phonophobia		4(4, 8)
nausea/vomiting		1(1, 2)
neck tightness		68 (21,9)*
deterioration on exertion		7(8, 4)
Numl	er of days/months with analgesic t	therapy
	10-14	2 (2,4)
	15-19	42 (50,6)*
	20-24	23 (27,7)*
	25-29	16 (19,3)
	Type of symptomatic therapy	i.
Combined		40 (48,2)*
Metamizole		2 (2,4)
NSAIDs		34 (41,0)*
Paracetamol		3 (3,6)
Triptans		4 (4,8)
	Efficacy of symptomatic therapy	L.
yes / partial		18 (21,7)
No		65 (78,3)*
	Type of prophylactic therapy	
TCAs		69 (83,1)*
AED		2 (2,4)
other		12 (14,5)
	Non-Pharmacological therapy	
Yes		1 (1,2)
No		81 (98,8)*
HIT		65,4±5,5
* Hi square test, p<0.05.	1	, ,

Table 2: Clinical Features of MOH

The correlation analysis revealed only the association between the duration of the previous headache and the effect of MOH on daily life activities (HIT) (r=0.327, p=0.003).

Discussion

The results of our research confirm the results of earlier studies (1, 3, 11 - 14) that MOH occurs mainly in middle age and more often in women.

Interestingly, the largest percentage of patients in our study did not smoke or consume alcohol, while the majority of them were physically inactive and regularly consumed caffeinated beverages. In the literature reviewed, no significant studies were found, the results of which could be useful for comparison with the results obtained here. A study examining various risk factors for MOH revealed a correlation between physical inactivity and smoking and MOH onset [9]. Our results are partially consistent with those findings. Studies examining the characteristics of patients who have never had

a headache in their lives have shown that they were men of younger ages who regularly consumed alcoholic beverages (unpublished results). Our findings on the inverse relationship between alcohol consumption and the occurrence of MOH are consistent with these findings. Although most patients with MOH do not have comorbidities, the incidence of cardiovascular, bone and joint diseases was significantly greater than that of other comorbid conditions. The reasons for these findings should be considered because arterial hypertension and spondylosis of the cervical spine (the most common diagnoses in these patients) are the most common causes of secondary chronic headaches [1]. This headache, due to its chronic course, poses a risk of developing MOH. Previous studies have shown an association between disease involving the osseous joint system and MOH [9].

Although MOH can arise from all primary headaches, MOH is most often caused by the transformation of chronic migraine [1]. The results of our study showed that the highest percentage of patients with MOH previously had chronic migraines. Our research has shown that MOH is most often caused by the excessive use of combined and ordinary analgesics (NSAIDs). The risk of MOH has previously been shown to be lower with triptans and ergotamine use than with NSAIDs [10]. Although they used prophylactic therapy, for the most part, patients with MOH did not achieve satisfactory therapeutic efficacy. The high values obtained here on the scale of assessment of the impact of headaches on daily life activities are in agreement with the results of previous studies [11].

Research devoted to examining the pathophysiological mechanisms of MOH has shown morphometric and functional changes in regions crucial for the modulation of nociception (periaqueductal gray matter, posterior cingular cortex, hippocampus, thalamus, orbitofrontal cortex, anterior cingular gyrus, insula, and precuneus) [12,13]. These structures are involved in nociception modulation, cognition, and affective and addictive behavior. The mechanisms of central sensitization (hyperexcitability of pain modulation pathways) are considered crucial in the pathophysiology of MOH [1]. Bilateral, moderately severe pain accompanied by tightness in the neck and shoulders and the absence of so-called migraine features of MOH in our study (even though it is largely due to the transformation of previous chronic migraine) support the previously assumed mechanisms of central sensitization.

Conclusion

MOH results from the transformation of a previous chronic headache due to overuse of analgesic therapy.

It affects the middle-aged population, mostly women, and affects all aspectso f the patient's life. Despite defined therapeutic strategies for treatment, prevention of MOH is considered inviolable (patient education, limiting the use of analgesic therapy, especially in the at-risk group of patients (middle-aged women with chronic migraine)). However, further investigations of other potentially modifiable risk factors for the development of MOH and its course and outcome after certain therapeutic interventions are necessary [14-17].

Declarations

Ethics Approval and Consent to Participate

- Consent for Publication: Yes.
- **Author Contributions:** SLJ made the initial plan and wrote the abstract, the introduction, and all the sections. RAD and ST participated in the planning of the Manuscript, the litera- ture list, and wrote the sections about medication overuse headache pathology.
- Funding: None.
- Data Availability Statement: On request.
- Acknowledgments: Marina Ljubisavljevic for her support in this paper revi sion.
- Conflicts of Interest: The authors have no conflicts of interest to declare.

References

- 1. Parisi, P., Belcastro, V., Verrotti, A., Striano, P., & Trenitè, D. K. N. (2018). "Ictal epileptic headache" and the revised International Headache Classification (ICHD-3) published in Cephalalgia 2018, vol. 38 (1) 1–211: Not just a matter of definition. *Epilepsy & Behavior*, *87*, 243-245.
- 2. Westergaard, M. L., Hansen, E. H., Glümer, C., Olesen, J., & Jensen, R. H. (2014). Definitions of medicationoveruse headache in population-based studies and their implications on prevalence estimates: a systematic review. *Cephalalgia*, 34(6), 409-425.
- 3. Okada-Ogawa, A., Porreca, F., & Meng, I. D. (2009). Sustained morphine-induced sensitization and loss of diffuse noxious inhibitory controls in dura-sensitive medullary dorsal horn neurons. *Journal of Neuroscience*, 29(50), 15828-15835.
- 4. Srikiatkhachorn, A., Tarasub, N., & Govitrapong, P. (2000). Effect of chronic analgesic exposure on the central serotonin system: a possible mechanism of analgesic abuse headache. *Headache: The Journal of Head and Face Pain, 40*(5), 343-350.
- 5. Tohyama, Y., Yamane, F., Merid, M. F., & Diksic, M. (2001). Effects of selective 5-HT1A receptor antagonists on regional serotonin synthesis in the rat brain: an autoradiographic study with a-[14C] methyl-L-tryptophan. *European Neuropsychopharmacology*, *11*(3), 193-202.
- 6. Dobson, C. F., Tohyama, Y., Diksic, M., & Hamel, E. (2004). Effects of acute or chronic administration of anti-migraine drugs sumatriptan and zolmitriptan on serotonin synthesis in the rat brain. *Cephalalgia*, 24(1), 2-11.
- 7. Evers, S., & Jensen, R. (2011). Treatment of medication overuse headache-guideline of the EFNS headache panel.

European Journal of Neurology, 18(9), 1115-1121.

- de Goffau, M. J., Klaver, A. R., Willemsen, M. G., Bindels, P. J., & Verhagen, A. P. (2017). The effectiveness of treatments for patients with medication overuse headache: a systematic review and meta-analysis. *The Journal* of *Pain*, 18(6), 615-627.
- Hagen, K., Albretsen, C., Vilming, S. T., Salvesen, R., Grønning, M., Helde, G., ... & Stovner, L. J. (2011). A 4-year follow-up of patients with medication-overuse headache previously included in a randomized multicentre study. *The journal of headache and pain*, *12*, 315-322.
- de Goffau, M. J., Klaver, A. R., Willemsen, M. G., Bindels, P. J., & Verhagen, A. P. (2017). The effectiveness of treatments for patients with medication overuse headache: a systematic review and meta-analysis. *The Journal* of Pain, 18(6), 615-627.
- 11. Mathew, N. T., Stubits, E., & Nigam, M. P. (1982). Transformation of episodic migraine into daily headache: analysis of factors. *Headache: The Journal of Head and Face Pain, 22*(2), 66-68.
- 12. Schwedt, T. J., & Chong, C. D. (2017). Medication overuse headache: pathophysiological insights from structural and functional brain MRI research. *Headache: The Journal of Head and Face Pain*, *57*(7), 1173-1178.
- 13. Lai, T. H., & Wang, S. J. (2018). Neuroimaging findings in patients with medication overuse headache. *Current pain and headache reports, 22*, 1-7.
- 14. Headache Classification Subcommittee of the International Headache Society (2004) The international classification of headache disorders: 2nd edition. *Cephalalgia 24*(Suppl 1):9–160.
- 15. Rapoport, A. M. (2008). Medication overuse headache: awareness, detection and treatment. CNS drugs, 22, 995-1004.
- 16. Ayzenberg, I., Katsarava, Z., Sborowski, A., Chernysh, M., Osipova, V., Tabeeva, G., ... & Lifting the Burden. (2012). The prevalence of primary headache disorders in Russia: a countrywide survey. *Cephalalgia*, *32*(5), 373-381.
- 17. Linde, M., Gustavsson, A., Stovner, L. J., Steiner, T. J., Barré, J., Katsarava, Z., ... & Andrée, C. (2012). The cost of headache disorders in Europe: the Eurolight project. *European journal of neurology*, *19*(5), 703-711.