Journal of Theoretical, Experimental, and Applied Physics ISSN: 3070-6505

PRIME OPEN ACCESS

Volume 2, Issue 4

Research Article

Date of Submission: 25 August, 2026
Date of Acceptance: 01 October, 2026
Date of Publication: 23 October, 2026

Near-Mass Pressure Illustrated: A Curvature-Driven Mechanism for Spherical Singularity Formation

David Nothstine*

Independent Researcher, Panama City, Florida, USA

***Corresponding Author:**
David Nothstine, Independent Researcher, Panama City, Florida, USA.

Citation: Nothstine, D. (2026). Near-Mass Pressure Illustrated: A Curvature-Driven Mechanism for Spherical Singularity Formation. *J Theor Exp Appl Phys, 2*(4), 01-21.

Abstract

This work proposes a curvature-driven mechanism for the formation of spherical singularities within the Deep Liquid Metallic Hydrogen (LMH) realm. In this framework, singularities do not originate from stellar collapse but arise through internal gravitational processes occurring under extreme pressure regimes deep within the universe.

At sufficiently high pressures, the LMH medium approaches a state described here as near-mass pressure, where stress–energy density and spacetime curvature approach critical thresholds. Within this metastable environment, localized instability nodes form where pressure and curvature gradients can no longer remain evenly distributed.

When these nodes exceed critical density, curvature, and geometric convergence thresholds, spacetime reorganizes into a confined spherical curvature structure. This transition represents a curvature-driven phase change in which distributed gravitational fields become self-reinforcing and form finite spherical singularities.

Over time, these singularities accumulate energy within their confinement geometry. When the stability limit of the curvature structure is exceeded, confinement failure occurs and the singularity releases its stored energy as an expanding spherical or helical curvature wavefront.

Within this cosmological framework, the interaction of successive singularity bursts produces overlapping spherical or helical wavefronts that shape the large-scale architecture of the universe. Filamentary structures, galaxy clusters, and cosmic voids may therefore represent the long-term geometric consequences of interacting curvature waves.

This model provides a structural mechanism linking deep-realm singularity formation to the emergence of large-scale cosmic structure.

Keywords: Near-Mass Pressure, Spacetime Curvature, Singularity Formation, Gravitational Convergence, Stress-Energy, Anisotropic Convergence, Curvature Locking, Cosmic Structure

Introduction

The mathematical divergence traditionally associated with singularities may reflect the limits of the equations used to describe a particular physical regime rather than the existence of an actual infinite point. This situation is analogous to attempting to use atmospheric equations to describe the dynamics of the deep ocean. The mathematics fails not because the ocean becomes infinite, but because the equations were never designed to operate under those conditions.

In a similar way, the divergence encountered in gravitational equations may signal the transition of spacetime into a different pressure and curvature regime rather than the breakdown of physical structure itself. Within the framework developed in earlier work, singularities arise within a deeper region of the universe described as the Deep Liquid Metallic Hydrogen (LMH) realm [1].

The purpose of this work is to describe the mechanism by which extreme pressure within this deep realm produces a curvature-driven phase transition—reaching what is here termed as the near-mass pressure threshold—that leads to the formation of a finite spherical singularity.

In this work I am not proposing a completely separate reality governed by unrelated laws. Rather, I suggest that the same underlying physical system is being pushed into conditions so extreme that the mathematical descriptions commonly used in our realm begin to diverge.

In the framework of General Relativity, mathematical divergence already appears when density becomes extremely large, curvature becomes extremely strong, and spacetime intervals approach limiting behavior. Under these conditions the equations begin producing infinities. Physicists generally interpret this not as nature itself becoming infinite, but as an indication that the mathematical model has reached the limit of its applicability. In other words, the physical regime has changed [2].

This pattern appears throughout physics. Ordinary equations describe ordinary conditions, while extreme regimes often require new descriptions. Mathematical divergence frequently signals a transition point between physical regimes. Examples include the transition from classical mechanics to quantum mechanics, laminar fluid flow to turbulence, phase transitions between states of matter, and the expected transition from classical General Relativity to a quantum-gravity regime.

In each case the mathematics ceases to behave smoothly because the assumptions underlying the model no longer hold.

Within the framework proposed here, the deep realm represents such a regime. It is a domain in which the conditions of pressure and curvature become so extreme that the mathematical descriptions commonly used for spacetime behavior in our observable realm no longer adequately describe the system.

The deep realm is not elsewhere.

It is the same physical continuum operating beyond the stability range of the observable spacetime behavior in the Relative Quantum Realm we live in.

Mechanism Chain of Near-Mass Pressure Singularity Formation
The sequence of processes proposed in this work can be summarized as the following mechanism chain.

Deep LMH Equilibrium
The Deep Liquid Metallic Hydrogen realm exists in a metastable state where pressure, density, and curvature remain balanced.

↓

Instability Node Formation
Small fluctuations produce localized regions where pressure and curvature gradients exceed the equilibrium capacity of the surrounding LMH medium.

↓

Near-Mass Pressure Threshold
Stress–energy density and curvature approach critical values.
The LMH medium can no longer distribute curvature smoothly.

↓

Curvature Convergence
Gravitational curvature begins to converge toward the instability node.

↓

Curvature Locking
Curvature becomes self-reinforcing. Spacetime reorganizes into a confined spherical curvature structure.

↓

Finite Spherical Singularity
A stable spherical singularity forms. Curvature becomes self-contained and dynamically stable within the structure.

↓

Energy Accumulation
Energy and curvature continue to build within the singularity's confinement geometry over time.

↓

Confinement Failure
When the stability limit of the curvature structure is exceeded, the singularity can no longer maintain confinement.

↓

Singularity Burst
The singularity releases stored energy as an expanding spherical or helical curvature wavefront.

Return to the Deep LMH realm and closure of the gravity well.

↓

Cosmic Structure Formation

Interactions between successive curvature wavefronts generate filamentary structures, galaxy clusters, and cosmic voids.

Core Equations of the Near-Mass Formation Model

Effective Gravitating Density

$\rho_{grav} = \rho + 3P/c^2$

Where

- ρ = mass density
- P = pressure

Near-mass pressure occurs when

$\rho_{grav} \geq \rho_c$

Curvature Indicators

Ricci Scalar

$R = -(8\pi G/c^4)T$

Kretschmann Scalar

$K = R_{abcd} R^{abcd}$

Near-mass curvature condition

$K \geq K_c$

Geometric Convergence Parameter

$C = r_n / L_{curv}$

Where

- r_n = radius of LMH instability node
- L_{curv} = curvature length scale

Curvature locking occurs when

$C \geq C_{th}$

Singularity Formation Condition

Spherical singularity formation occurs when

$\rho_{grav} \geq \rho_c, \quad K \geq K_c, \quad C \geq C_{th}$

These thresholds indicate the transition from distributed curvature to a self-contained spherical gravitational structure.

Singularity Burst Expansion

Following confinement failure, curvature wave fronts propagate outward approximately as

$r(t) = r_0 + vt$

Where

- r_0 = initial singularity radius
- v = expansion velocity

Curvature strength decreases with distance

$K(r) \propto 1/r^2$

The Deep LMH Realm as a Curvature-Saturated Medium
In the framework of the Three Realms theory, singularities are not created by stellar collapse from the relative realm. Instead, they arise within a deeper regime of the universe: the Deep Liquid Metallic Hydrogen Realm. D.L.M.H.R.

This realm exists under immense pressures and energy densities where matter and spacetime curvature are tightly coupled. At such depths, the medium is not empty or smooth. It is a compressed gravitational environment in which curvature already permeates the structure of spacetime.

Unlike ordinary astrophysical environments, the LMH realm exists near a state of metastable equilibrium. Pressure, density, and curvature remain balanced across large regions, yet this equilibrium is not perfectly stable. Small fluctuations inevitably arise.

These fluctuations manifest as localized instability nodes.

Within such nodes the stress–energy density of the medium begins to exceed the equilibrium capacity of the surrounding LMH field. As a result, curvature can no longer remain evenly distributed. Instead, spacetime begins to converge around the instability node.

The near-mass pressure effect represents the mechanism by which the Deep LMH realm stabilizes such instability nodes. When pressure and curvature exceed equilibrium limits, gravitational convergence reorganizes the region into a confined spherical singularity structure.

This transition defines the condition described in this work as near-mass pressure.

Stress–Energy and the Approach to Near-Mass Pressure
In general relativity, gravitational curvature is determined by the Einstein field equations [3].

$$G_{\mu\nu} = (8\pi G/c^4)\, T_{\mu\nu}$$

where
- $G_{\mu\nu}$ is the Einstein tensor, describing spacetime curvature.
- $T_{\mu\nu}$ is the stress–energy tensor, describing the distribution and flow of energy, momentum, pressure, and matter.
- G is the gravitational constant.
- c is the speed of light in vacuum.

The stress–energy tensor includes contributions not only from mass density but also from pressure and internal energy.

For a relativistic fluid the effective gravitating density can be approximated by

$$\rho_{grav} = \rho + 3P/c^2$$

where
- ρ is mass density
- P is pressure

Under extreme compression, pressure contributes significantly to the gravitational field.
In the Deep LMH realm the pressure becomes so large that the effective gravitating density approaches a critical threshold

$$\rho_{grav} \geq \rho_c$$

where ρ_c represents the near-mass critical density.

At this point the LMH medium enters a transitional regime where curvature can begin to organize into a gravitationally autonomous structure.

Curvature Indicators and the Onset of Convergence
The presence of strong gravitational curvature can be measured using curvature invariants.

One useful quantity is the **Ricci Scalar**

$$R = -(8\pi G/c^4)T$$

where T is the trace of the stress–energy tensor.

Another measure is the **Kretschmann Scalar**

$K = R_{abcd} R^{abcd}$

which represents the total magnitude of spacetime curvature.

In the Deep LMH regime, near-mass pressure corresponds to a condition in which curvature approaches a critical value:

$K \geq K_c$

When this threshold is reached, curvature gradients intensify around instability nodes.

Spacetime no longer behaves as a smoothly distributed field.

Instead, curvature begins to focus and converge.

Geometric Convergence and the Instability Node

The instability node acts as a geometric anchor for this convergence.

- r_n represents the characteristic radius of the LMH instability node
- L_{curv} represents the local curvature length scale

A geometric convergence parameter can be defined as

$C = r_n / L_{curv}$

When curvature variations across the node become comparable to the node's scale,

$C \geq C_{th}$

the surrounding geometry can no longer remain diffuse.

Instead, curvature begins to fold inward toward the node.

This process marks the beginning of a geometric phase transition.

The concepts explored throughout Near-Mass Pressure involve geometric behaviors and evolving organizational structures that are often difficult to fully communicate through mathematics and descriptive language alone.

These upcoming illustrations had been developed as a visual extension of the framework, intended to explore how curvature organization, instability formation, anisotropic convergence, and recursive structural evolution can emerge within spacetime approaching near-mass pressure conditions.

Beginning with early multi-axis curvature development prior to prenode formation. Then moving to the original conceptual prenode sketch and progressing through orthogonal sectional analyses, layered convergence structures, and stratified knot analogues, the illustrations attempt to provide spatial intuition for the geometric processes described throughout the framework.

Together, these figures form a progressive visual interpretation of evolving curvature systems, where localized instability regions may organize dynamically through directional pathways, layered reinforcement, and multiple interacting axes of convergence rather than through idealized isotropic collapse.

Rather than functioning solely as artistic representations, the illustrations serve as conceptual geometry tools designed to externalize internal spatial reasoning into observable structural forms. In doing so, they provide a parallel visual language accompanying the theoretical and mathematical foundations presented throughout Near-Mass Pressure.

This illustrates how multiple localized regions begin operating simultaneously around a developing central instability region, producing anisotropic distortions and the first emergence of distinct organizational axes.

At this stage, no stable central prenode is yet established. Instead, the converging curvature pathways begin concentrating toward a shared central region.

The innermost structures remain irregular and strongly directional, while successive outer layers spread and level the deformation across a broader curvature envelope. Within the Near-Mass Pressure framework, this figure represents the transitional geometry through which distributed curvature amplification, layered feedback, and multi-axis convergence begin organizing toward prenode formation.




Figure A: Early Multi-Axis Curvature Development Prior to Prenode Formation

In the planar view, the inner curvature remains irregular and strongly directional. Separate convergence zones form around the center, with each region contributing differently to the developing geometry. These competing but interconnected pathways begin concentrating curvature toward a shared central location.

The 90-degree cutaway reveals that this organization is not confined to a single plane. The emerging structure extends through orthogonal directions, with primary polar and equatorial axes developing together while secondary convergence regions form between them. This gives the early curvature field a three-dimensional, multi-axis character.

As the structure extends outward, successive curvature layers distribute the inner irregularities across a broader region. The outer layers therefore become progressively smoother and more level with distance, forming an enclosing feedback structure around the developing center.

Within the Near-Mass Pressure framework, the illustration suggests that asymmetry is not a defect in the geometry. It is a natural part of the dynamical process through which multiple convergence zones, organizational axes, and layered curvature feedback begin forming prior to prenode development.




Figure 1

"The shaded external layering represents the evolving curvature organization surrounding the prenode, while the labeled internal strands conceptually represent emerging pathways of stress, density, and curvature alignment within the structure.

The apparent disappearance of several shaded curvature lines near the north-end convergence region is not an error in the illustration, but rather an intentional representation of layered geometric overlap within the evolving prenode structure.

The curvature pathways remain connected, yet portions become visually obscured as stronger foreground curvature layers intensify and overlap during convergence.

This effect reflects the dynamic, three-dimensional nature of the prenode geometry, where organizational layers interact recursively rather than remaining isolated or uniformly visible. What emerges is the appearance of curvature feeding back into itself through layered reinforcement and directional concentration.

Different regions of the structure dominate visually depending on depth, orientation, and convergence intensity, producing partial concealment of underlying pathways.

This behavior supports the interpretation that evolving instability systems naturally develop anisotropic and self-reinforcing organizational structures, where internal and external curvature geometries become dynamically coupled during the approach toward near-mass pressure conditions.

This is the structure I envisioned within my mind and later sketched outward into visual form.

As we move progressively through the illustrations, a visual conception begins to emerge of the underlying geometric and organizational principles described throughout Near-Mass Pressure.

What may initially appear as isolated conceptual forms gradually develops into an interconnected structural interpretation of evolving curvature systems within spacetime.

Each illustration builds upon the previous one, revealing additional layers of organization, anisotropic behavior, recursive convergence, and directional curvature evolution. Together, the figures form a visual progression that mirrors the theoretical development of the framework itself.

These illustrations are therefore not intended merely as artistic representations, but as geometric interpretation tools designed to assist in visualizing processes that are otherwise difficult to fully communicate through mathematics and descriptive language alone.

Through layered curvature structures, orthogonal sectional views, recursive overlap, convergence ridges, channel-like pathways, and knot-like stratified organization, the figures attempt to translate abstract curvature dynamics into observable conceptual geometry.

As the sequence progresses, you are gradually introduced to the possibility that evolving instability systems within spacetime may organize themselves through directional pathways rather than through idealized isotropic collapse.

The illustrations increasingly suggest that localized curvature amplification can produce layered feedback structures, multiple organizational axes, anisotropic convergence zones, and progressively intensifying concentration regions approaching near-mass pressure conditions.

Ultimately, the illustrations represent an effort to externally translate an internal geometric vision into a structured visual form that can be observed, examined, and conceptually explored alongside the theoretical framework presented throughout Near-Mass Pressure.

Within the present framework, asymmetry is not regarded as an anomaly or deviation from an otherwise perfect geometric order, but rather as a fundamental and natural consequence of the dynamical evolution of physical systems.

While idealized symmetric models serve as valuable tools for mathematical simplicity and first-order approximations, real-world systems rarely sustain perfect symmetry once subject to dynamic processes.

Environmental influences constantly act upon these systems, leading to the spontaneous development of directional organization, localized concentration regions, anisotropic flow behaviors, and geometrically uneven structures over time.



PRENODE – ASYMMETRIC STRUCTURE WITH SOUTH CORE CONCENTRATION
Two Orthogonal Slices of the Same Structure (Quarter-Turn Difference)
PRE-NODE OVERVIEW
A transitional, non-spherical instability structure in which spacetime curvature, density, and pressure become organized into internal ridges and channels prior to node (singularity) confinement.
NORTH APEX CONVERGENCE ZONE
Narrow, highly compressed region where curvature, pressure, and density gradients converge toward incipient critical conditions.
RIDGES OF CURVATURE (ELONGATED)
Elongated, load-bearing structures of raised curvature that align with dominant convergence flow and carry the highest structural stress.
INTERNAL CHANNELS
Lower-curvature pathways between ridges that guide the flow of stress, density, and geometric information along the axis.
EQUATORIAL TRANSITION ZONE
Region of reorganization and redistribution. Internal stresses are redistributed as flow begins to focus more strongly toward the south core.
LAMINAR BOUNDARY
Outer envelope containing the prenode structure. Defines the transitional instability region and separates the organized interior from the exterior spacetime environment.
SOUTH CORE CONCENTRATION
Asymmetric core at the south end where curvature, density, and pressure reach their maximum concentration. All internal flow converges into this core.
SLICE A – LONGITUDINAL VIEW (A–A')
(North Apex → South Core)
CURVATURE FIELD LINES – DRAWN OUT TO THE EXTERIOR
THE SAME PRE-NODE STRUCTURE VIEWED FROM TWO ORTHOGONAL PLANES
A quarter-turn (90° rotation) between slices reveals the full three-dimensional anisotropic geometry.
A
B
B'
A'
Curvature is concentrated at the south core and gradually drawn out along ridges and channels, extending into the surrounding field. The field lines reflect anisotropic flow and structural influence reaching far beyond the prenode boundary.
SLICE B – ORTHOGONAL VIEW (B–B')
(Rotated 90° from Slice A)
PRE-NODE OVERVIEW
A transitional, non-spherical instability structure in which spacetime curvature, density, and pressure become organized into internal ridges and channels prior to node (singularity) confinement.
NORTH APEX CONVERGENCE ZONE
Narrow, highly compressed region where curvature, pressure, and density gradients converge toward incipient critical conditions.
RIDGES OF CURVATURE (ELONGATED)
Elongated, load-bearing structures of raised curvature that align with dominant convergence flow and carry the highest structural stress.
INTERNAL CHANNELS
Lower-curvature pathways between ridges that guide the flow of stress, density, and geometric information throughout the structure.
EQUATORIAL TRANSITION ZONE
Region of reorganization and redistribution. Internal stresses are redistributed as flow begins to focus more strongly toward the south core.
LAMINAR BOUNDARY
Outer envelope containing the prenode structure. Defines the transitional instability region and separates the organized interior from the exterior spacetime environment.
SOUTH CORE CONCENTRATION
Asymmetric core at the south end where curvature, density, and pressure reach their maximum concentration. All internal flow converges into this core.
INTERPRETATION
The prenode is an anisotropic, organized structure. Its interior contains an asymmetric core of maximum concentration at the south end. Curvature, density, and pressure are organized into ridges and channels that draw flow outward, influencing the surrounding spacetime field. The structure is not yet a singularity, but a transitional instability region approaching node confinement.
CROSS-SECTION NEAR SOUTH CORE (VIEW A–A' NEAR CORE)
Internal ridges encircle and feed the south core.
Channels between ridges guide flow inward.
Strong curvature gradients compress inward toward the asymmetric core.
This cross-section shows the organized compression pattern near the core region.
RELATIONSHIP
Slices A and B are orthogonal views (quarter-turn difference) of the same 3D structure. Together they reveal the full anisotropic geometry.
CROSS-SECTION NEAR SOUTH CORE (VIEW B–B' NEAR CORE)
Internal ridges encircle and feed the south core.
Channels between ridges guide flow inward.
Strong curvature gradients compress inward toward the asymmetric core.
This cross-section shows the organized compression pattern near the core region.
FLOW DIRECTION SUMMARY
All internal pathways and curvature ridges guide stress, density, and geometric flow from the north apex region toward the south core concentration.
North Apex → South Core
Curvature is concentrated at one end and drawn out to the field.
COLOR GUIDE:
Low Density / Low Curvature
Moderate Density / Moderate Curvature
Elevated Density / Elevated Curvature
High Density / High Curvature
Maximum Density / Maximum Curvature
Direction of Curvature Flow

Figure 2




Figure 3

Multiple Organizational Axes and Anisotropic Prenode Geometry

An important implication emerging from the present framework is the possibility that the prenode structure possesses multiple organizational axes simultaneously. This interpretation arises from examining orthogonal slice geometries through the same transitional instability structure, where quarter-turn rotational views reveal differing internal ridge alignments, compression pathways, and curvature-channel organization. Such behavior suggests that the prenode is not rotationally uniform or isotropic, but instead contains internally organized directional structures shaped by evolving curvature dynamics and anisotropic stress redistribution.




Figure 4

In highly symmetric systems, rotational slices separated by 90° would produce nearly identical internal geometries. However, within the present prenode framework, differing slice orientations expose variations in ridge concentration, channel morphology, curvature gradients, and flow organization. This indicates that the structure could easily include multiple interacting axes of convergence and stress transport operating simultaneously within the evolving instability region.

This type of anisotropic organizational behavior is not without precedent in natural systems. Comparable phenomena appear in:

- Fluid Instabilities
- Magnetic Confinement Systems
- Plasma Organization
- Crystal Growth
- and stress-field redistribution within structured materials.

In such systems, organization frequently emerges through directional pathways rather than perfectly uniform collapse. Internal flow channels, ridges, vortices, and stress-alignment structures develop dynamically as energy, pressure, or curvature redistribute throughout the medium. The resulting geometries are often highly organized, yet non-uniform and axis-dependent.

Within the present framework, the prenode increasingly appears less like a static object and more like an evolving organized field structure embedded within spacetime itself. Curvature, density, and near-mass pressure may propagate preferentially along structured ridges and channels that guide convergence toward localized concentration regions. The structure therefore behaves as a transitional geometric topology in which multiple directional organizational processes coexist prior to confinement stabilization.

This interpretation may have important implications for understanding the emergence of instability nodes and the transition toward spherical singularity formation. Rather than arising from perfectly isotropic collapse, confinement may emerge from the progressive interaction and tightening of anisotropic curvature pathways that eventually reorganize into a more stable confined geometry through curvature locking and near-mass pressure concentration.

Consequently, the prenode may represent a dynamically evolving spacetime-organizing structure whose geometry reflects the directional redistribution of curvature, pressure, and stress across multiple interacting axes. Such behavior would naturally produce differing internal morphologies depending upon observational orientation or sectional geometry, consistent with the orthogonal slice behavior illustrated within the present framework.




Figure 5

You can see in this picture how the surrounding layers concentrate around the core. This serves as a natural two-dimensional representation of the conceptual idea proposed in this work: that curvature may progressively organize and intensify around a localized instability node.

Procedural wood-knot generation provides a useful structural analogy for understanding how asymmetry and localized organization may naturally emerge within evolving curvature systems. In wood formation, growth occurs through layered concentric expansion around a central axis, yet the presence of a branch or localized disturbance produces a knot—a region where the surrounding grain becomes warped, compressed, twisted, and directionally reorganized around a localized attractor.

A simplified radial growth field may be represented by:
$r = \sqrt{(x^2 + y^2)}$

with layered structure approximated through a periodic function such as:

$f(r) = \sin(r\omega)$

where ω represents the spatial frequency of the layered growth pattern. In an idealized system, such layers would remain smooth and symmetric. However, the introduction of a localized attractor at position:

$\vec{C} = (c_x, c_y, c_z)$

distorts the surrounding coordinate field. The distance from a point $\vec{P}$ to the attractor is:

$d = \|\vec{P} - \vec{C}\|$

and the strength of the local distortion may be represented by a falloff function:

$w = \max(0, 1 - d/R)^2$

where R defines the effective radius of influence around the attractor. The distorted coordinate field may then be expressed schematically as:

$\vec{P}' = \vec{P} + w\vec{D}(\vec{P})$

where $\vec{D}(\vec{P})$ represents the local displacement or twisting field generated by the attractor.

A remarkably similar behavior may occur within evolving spacetime geometries approaching near-mass pressure conditions. In this interpretation, the prenode functions analogously to a curvature attractor embedded within stratified spacetime, where local curvature amplification and stress-energy redistribution distort surrounding geometric pathways into organized ridges, channels, and convergence structures. The curvature concentration may be represented through a curvature invariant such as:

$K(r) = R_{abcd} R^{abcd}$

with localized increases in K(r) corresponding to intensified curvature around the developing instability node.

As curvature gradients intensify,

$\nabla K \neq 0$

spacetime may no longer evolve through idealized isotropic collapse, but instead through warped directional organization resembling layered knot formation within natural growth systems. The resulting geometry becomes increasingly anisotropic, producing localized curvature concentration zones and multiple interacting axes of convergence that ultimately contribute to instability growth, curvature locking, and confinement emergence.

Phase Transition and Singularity Formation
The formation of a spherical singularity occurs when the following conditions are simultaneously satisfied:

$\rho_{grav} \geq \rho_c, \quad K \geq K_c, \quad C \geq C_{th}$

At this threshold the LMH medium undergoes a curvature-driven phase transition. Distributed curvature reorganizes into a confined region of spacetime geometry, and the gravitational field becomes self-reinforcing, locking around the instability node.

The result is the emergence of a finite spherical singularity.

Unlike classical point-singularity descriptions, this structure is not a mathematical divergence but a stable geometric configuration in which curvature becomes concentrated and self-sustaining. In this framework, singularity formation represents the mechanism through which equilibrium is restored.

The structure that emerges from this stabilization process is spherical. A sphere provides the most uniform geometry through which pressure and curvature can be distributed around a confined region. By concentrating curvature equally in all directions, the spherical form offers the most stable configuration capable of containing the instability node.

This outcome is consistent with broader patterns observed throughout the universe. Across many scales of nature, spherical structures arise wherever forces act uniformly toward a central region. From stars and planets to gravitational systems, the sphere repeatedly appears as the natural geometry of equilibrium and confinement.

For this reason, the stabilization of instability nodes within the Deep LMH realm resolves into the formation of spherical singularities.

The Knot: Curvature Around a Node

A Geometric Analogue for Anisotropic Curvature Organization Within a Stratified Medium

This figure illustrates a conceptual geometric analogue for how localized curvature organization may emerge within a layered spacetime environment approaching near-mass pressure conditions. Similar to the way wood grain bends, compresses, twists, and reorganizes around a knot, stratified curvature layers may progressively distort around a developing instability region embedded within a curvature-saturated medium.

The knot-like structure represents a localized prenode concentration region where curvature, pressure, density, and directional stress redistribution become increasingly organized through recursive convergence and layered reinforcement. Rather than evolving through idealized isotropic collapse, the surrounding geometry appears to reorganize through anisotropic pathways that guide convergence toward localized regions of intensified curvature concentration.

The layered wrapping behavior shown throughout the structure serves as an analogue for evolving curvature organization within spacetime. As local curvature gradients intensify, surrounding geometric pathways may bend inward, overlap, compress, and recursively reinforce one another, producing organized ridges, channels, and spiral-like convergence structures. This process reflects the emergence of directional curvature organization rather than smooth symmetric collapse.




Figure 6

The three-dimensional conceptual view demonstrates how layered stratification may support multiple interacting organizational pathways simultaneously. Curved inflow trajectories, rotational organization, and recursive directional convergence suggest that evolving instability systems may naturally develop anisotropic internal geometries with multiple interacting axes of concentration.

The cross-sectional interpretation further illustrates how surrounding layers may progressively narrow and reorganize toward localized convergence regions. Internal curvature pathways become increasingly compressed as directional stress redistribution intensifies near the prenode concentration zone.

Likewise, the top-down view emphasizes the possibility of spiral and rotational organizational behavior emerging within evolving curvature systems. Rather than representing simple circular motion, the spiral structure symbolizes recursive geometric reinforcement and the directional redistribution of curvature through layered organizational pathways.

The formation sequence shown at the bottom of the figure is best interpreted not as discrete object formation, but as a progressive organizational evolution:

- Stratified Medium
- Local Curvature Distortion
- Directional Curvature Organization
- Recursive Convergence Amplification
- Prenode Concentration Region

Within this interpretation, the prenode behaves less like a static object and more like an evolving geometric instability topology embedded within stratified spacetime. Curvature gradients progressively organize surrounding layers into anisotropic convergence pathways that intensify through recursive feedback and directional reinforcement during the approach toward near-mass pressure conditions.

The knot analogy therefore serves not merely as a visual metaphor, but as a structural interpretation tool for understanding how layered curvature organization, recursive overlap, anisotropic convergence, and localized instability concentration may naturally emerge within evolving spacetime geometries.

The concepts explored throughout Near-Mass Pressure often emerged within my mind first as evolving geometric structures, layered organizational patterns, and dynamic curvature relationships long before they could be translated into mathematical language or formal description. Through these illustrations, I have attempted to externalize those internal geometric visions into forms that can be visually observed, examined, and conceptually explored. Each figure represents not merely an artistic rendering, but an effort to communicate the spatial intuition, recursive organization, anisotropic behavior, and evolving curvature dynamics that form the foundation of the framework presented throughout this work.

Relation to the Geometric Capture of Light

This process mirrors a principle described elsewhere in my work as the quantum trapping, or geometric capture of light.

Within a spherical singularity, photons continue to move at the speed of light along geodesic paths. However, the curvature of spacetime bends those paths until they become confined within the geometry, effectively circling a geometric center of extreme curvature.

The same principle governs singularity formation itself. When curvature exceeds a critical threshold in the Deep LMH realm, spacetime becomes capable of trapping its own geometry. Curvature no longer disperses outward but instead organizes into a stable spherical structure.

Thus, the formation of a singularity and the confinement of light within it both arise from the same fundamental principle: the geometric dominance of curvature over motion.

Instability Nucleation in the Deep LMH Field

Within the Deep Liquid Metallic Hydrogen (LMH) realm, equilibrium is maintained by the immense pressures that dominate this region of the universe. At such depths the medium behaves as a highly compressed gravitational field in which curvature, energy density, and pressure remain tightly coupled.

Despite the apparent stability of this environment, perfect equilibrium cannot be sustained indefinitely. Under extreme pressure regimes, even small perturbations in density or curvature can grow into localized regions of instability.

These regions represent the nucleation points of singularity formation.

Instability nucleation occurs when local fluctuations in pressure or density create gradients that cannot be smoothly redistributed through the surrounding LMH medium. When these gradients exceed a critical threshold, the stress–energy distribution begins to reorganize.

This condition may be expressed through a density gradient criterion

$|\nabla\rho| \geq (\nabla\rho)_c$

or a pressure gradient condition

$|\nabla P| \geq (\nabla P)_c$

where
- $(\nabla\rho)_c$ is the critical density gradient
- $(\nabla P)_c$ is the critical pressure gradient

When these thresholds are exceeded, the local region becomes dynamically unstable. The LMH medium can no longer distribute the gravitational load evenly across the surrounding volume.

Instead, curvature begins to converge toward the instability node.

This convergence marks the beginning of gravitational nucleation, where curvature gradients amplify and begin to fold inward toward the unstable region.

The instability node therefore becomes the geometric seed around which a spherical singularity can form and stabilize the system once again.

Curvature Locking and the Emergence of the Spherical Singularity

As curvature converges around the instability node, the surrounding geometry begins to reorganize. The local gravitational field intensifies as the stress–energy density approaches the near mass threshold described earlier.

During this stage, spacetime curvature transitions from a distributed field into a confined structure. This process can be described as curvature locking.

Curvature locking occurs when gravitational convergence becomes self-reinforcing. In this regime, curvature gradients feed back into the stress–energy distribution, further intensifying the gravitational field around the node.

Mathematically, this condition can be expressed when the convergence parameter C approaches unity:

$C = r_n / L_{curv} \rightarrow 1$

where r_n represents the characteristic radius of the instability node and L_{curv} is the local curvature scale of spacetime.

At this threshold, the curvature scale of spacetime becomes comparable to the physical scale of the instability node itself. The geometry can no longer remain diffused. Instead, spacetime reorganizes into a closed spherical curvature structure.

This structure represents the birth of a finite spherical singularity.

The singularity is therefore not an infinitely dense point but a geometric region in which curvature has become self-contained and dynamically stable. Within this spherical region, gravitational confinement becomes dominant. Motion within the singularity becomes governed primarily by the internal curvature geometry rather than by the surrounding LMH medium.

The gravitational structure produced by the presence of the singularity within the Relative Quantum Realm—described in earlier work, as the gravity bubble—corresponds observationally to what is commonly interpreted as a black hole [4].

As the newly formed singularity begins to acquire angular momentum as it moves through the gravity well, rotational dynamics develop within the confined curvature structure. Rotation organizes surrounding curvature and energy flows, generating increasingly powerful magnetic fields.

As these fields intensify, their influence begins to extend outward from the singularity's confined region into the surrounding spacetime. At this stage, the gravitational influence of the singularity becomes externally visible.

The result is the formation of the gravity bubble within our realm — the structure commonly identified as a black hole. The gravity bubble is not the singularity itself but the gravitational projection of the singularity into surrounding spacetime.

As angular momentum increases, both gravitational amplification and magnetic field strength increase. Rotation therefore plays a critical role in coupling the deep singularity structure to observable astrophysical phenomena within our realm.

Curvature locking refers to the self-reinforcing gravitational convergence that occurs when the curvature scale of spacetime approaches the physical scale of an instability node.

Confinement Dynamics and the Storage of Mass-Energy and Curvature

Once the spherical singularity forms, it acts as a gravitational confinement structure. Energy and matter entering the singularity become trapped within the curved spacetime geometry in a state of near-stasis at the quantum level. Rather than being destroyed, these constituents become incorporated into the internal gravitational configuration of the singularity.

The singularity therefore behaves as a reservoir of stored mass-energy and curvature.

Over time, continued accretion from the Relative Quantum Realm increases the mass-energy density within the singularity. As the internal mass-energy density rises, the curvature of the singularity intensifies further.

However, this process cannot continue indefinitely.

Every confinement structure possesses a stability limit.

As internal curvature and energy density increase in the Relative Quantum Realm (our universe), the singularity approaches a critical confinement threshold.

When this threshold is exceeded, the stability of the structure begins to weaken, as developed in earlier work [5].

Confinement Failure and the Singularity Burst

When the internal mass-energy of the singularity surpasses its structural stability limit, the confinement geometry can no longer maintain equilibrium.

This leads to a process described in earlier works as singularity burst mechanics.

During a burst event, the stored mass-energy and curvature contained within the singularity are suddenly released into the surrounding spacetime.

The release propagates outward as spherical or helical curvature wavefronts depending on rotation or non-rotating singularities, carrying matter, radiation, and energy into the surrounding cosmic environment.

These expanding wavefronts form the structural seeds of cosmic architecture.

In the framework described in earlier works, overlapping singularity bursts produce interacting spherical or helical wavefronts that sculpt the large-scale structure of the universe [4,5].

The cosmic web, galaxy clusters, and filamentary structures can therefore be interpreted as the long-term consequences of these interacting curvature waves.

Thus the life cycle of a singularity may be summarized as:

LMH instability nucleation
- Curvature convergence
- Spherical singularity formation
- Gravitational confinement
- Mass-energy accumulation
- Confinement failure
- Singularity burst
- Return to the Deep LMH realm and closure of the gravity well

This sequence forms the mechanical backbone of the cosmological processes described throughout these works.

Spherical and Helical Wavefront Propagation Following Singularity Burst
When the confinement geometry of a spherical singularity fails, the stored mass-energy and curvature within the singularity are released into the surrounding spacetime. This release does not occur as a random dispersal of matter or radiation. Instead, the geometry of the singularity determines the form of the expansion.

Because the singularity exists as a spherical curvature structure, the release of its stored mass/energy propagates outward in the form of Spherical or Helical wavefronts.

These wavefronts represent moving regions of enhanced curvature traveling through spacetime. Matter, radiation, and energy are carried within these expanding shells as the curvature propagates outward from the point of release approaching the speed of light c.

The propagation of these curvature waves can be approximated by a radial expansion relation

$r(t) = r_0 + vt$

where
- r(t) is the radius of the expanding wavefront
- r_0 is the initial radius of the singularity
- v represents the outward propagation velocity of the curvature disturbance

In many cases this velocity approaches relativistic limits due to the extreme L.M.H. expansion released during the burst event.

As the wavefronts expand, the curvature intensity decreases with distance from the origin. This reduction can be expressed as an approximate inverse square relationship

$K(r) \propto 1/r^2$

where K(r) represents the curvature magnitude at radius r.

This decreasing curvature gradient allows the expanding wavefront to gradually transition from an intensely curved gravitational disturbance into a broader region of distributed matter and energy.

Intersecting Wavefronts and the Emergence of Cosmic Structure
The universe described in the Genesis framework is not produced by a single burst event but by many singularity bursts occurring across Relative Quantum Realm over vast spans of time.

Each burst generates wavefronts expanding through spacetime. As these wavefronts propagate outward, they inevitably encounter the wavefronts produced by neighboring singularity bursts.

Where these expanding shells intersect, regions of constructive interaction occur.

At these intersection zones, curvature and matter densities combine, producing localized regions of enhanced gravitational attraction. These regions become the seeds of large-scale cosmic structures.

The intersection of multiple spherical wavefronts naturally produces the filamentary structures observed throughout the universe.

Matter flowing along intersecting curvature gradients becomes concentrated along elongated paths, forming the large cosmic filaments that connect galaxy clusters across intergalactic space.

This process can be visualized as the interaction of expanding spherical or helical shells within a vast three-dimensional medium.

The boundaries of these shells create zones of compression where the energy densities from multiple wavefronts reinforce one another.

These compression zones become the birthplaces of galaxies, clusters, and other cosmic structures.

Nested Wavefronts and Generational Structure (As Described in Genesis)
Not all singularity bursts occur simultaneously.

The deep realm operates across enormous spans of time, and bursts may occur in successive generations. As a result, newer wavefronts propagate through a universe already shaped by earlier bursts.

This creates a layered system of nested burst waves.

Older wavefronts expand to enormous scales, forming the largest cosmic voids and superstructures. Younger bursts occur within these regions, producing smaller wavefronts that interact with the existing curvature environment.

This generational layering naturally explains the presence of cosmic structures on many different scales.

Large voids and superclusters emerge from early large-scale bursts, while smaller galaxies and local structures arise from later, smaller bursts occurring within already established cosmic environments.

The Cosmic Web as a Fossil Record of Wave Intersections

Within this framework, the large-scale structure of the universe can be interpreted as the fossil record of interacting spherical wavefronts.

The cosmic web observed by modern astronomical surveys displays a network of filaments, clusters, and voids extending across billions of light-years.

In the Genesis model, these structures are not random gravitational accumulations but the long-term result of curvature waves intersecting and reinforcing one another over cosmic time [5].

Where wavefront intersections are strongest, clusters of galaxies form. Where wavefronts pass without strong interaction, large cosmic voids remain.

Thus the cosmic web represents the geometric memory of past singularity bursts.

The universe becomes a record of curvature interactions propagated through spacetime across immense epochs.

The Continuous Cycle of Singularity Formation and Burst

The processes described in this work reveal a continuous cycle operating within the cosmos.

Within the Deep LMH realm, instability nodes form under extreme pressure. These nodes develop into spherical singularities through curvature convergence.

Over time these singularities accumulate in our universe mass/energy and curvature until their confinement geometry fails. The resulting burst releases wavefronts that shape cosmic structure throughout the Relative Quantum Realm.

These wavefronts propagate outward, intersect, and generate the large-scale architecture of the universe.

Meanwhile, new instability nodes continue forming within the deep realm, producing new generations of singularities.

In this way the cosmos becomes a living dynamical system, continuously shaped by the creation, evolution, and bursting of spherical singularities.

The universe is therefore not a static structure nor the aftermath of a single primordial event but many primordial events over eons of time. It is an evolving architecture produced by the repeated interaction of gravitational structures emerging from the deep realm.

Observational Consequences of Singularity Burst Cosmology

As described in earlier work [5].

If the universe is shaped by successive generations of spherical singularity bursts, then the large-scale structure of the cosmos should preserve the geometric signatures of these events. Rather than emerging from a single primordial expansion, cosmic structure would reflect the cumulative effects of interacting curvature wavefronts propagating through spacetime.

Final Synthesis

Near-Mass Pressure and the Architecture of Cosmic Structure

The framework presented in this work describes a curvature-driven mechanism through which spherical singularities form within the Deep Liquid Metallic Hydrogen (LMH) realm. In this model the formation of singularities is not the result of stellar collapse but emerges from internal gravitational processes occurring within a curvature-saturated medium existing under extreme pressure regimes.

Within the Deep LMH realm, pressure, density, and spacetime curvature exist in a metastable equilibrium. Small fluctuations within this environment generate localized instability nodes where stress–energy density and curvature gradients can no longer remain evenly distributed. As pressure conditions intensify, the LMH medium approaches the regime described here as near-mass pressure, where curvature begins to dominate the behavior of the system.

When instability nodes exceed critical geometric and curvature thresholds, gravitational convergence becomes self-reinforcing. This process leads to curvature locking, in which spacetime reorganizes into a confined spherical curvature structure. The resulting configuration is a finite spherical singularity in which curvature becomes self-contained and dynamically stable.

Within this confinement geometry, mass–energy and curvature accumulate over time. The singularity functions as a storage structure in which gravitational energy is held within the locked curvature field. When the stability limits of this configuration are exceeded, the confinement structure fails and the singularity releases its stored energy in the form of an expanding curvature wavefront.

These singularity bursts propagate outward through spacetime as spherical or helical curvature waves. The interaction of successive bursts produces overlapping wavefronts that generate large-scale geometric patterns throughout the universe. Filamentary structures, galaxy clusters, and cosmic voids may therefore represent the long-term consequences of intersecting curvature waves generated by repeated singularity formation and release.

Within this framework, the cosmic web can be interpreted as a fossil record of intersecting curvature wavefronts produced over multiple generations of singularity bursts. Large-scale cosmic structure thus emerges not from a single primordial event but from an ongoing cycle of formation, confinement, and release occurring within the Deep LMH medium.

The concept of near-mass pressure therefore provides a structural mechanism linking deep-realm gravitational dynamics to the emergence of observable cosmic architecture. Rather than representing endpoints of gravitational collapse, singularities become active components within a continuous cosmological cycle.

This perspective reframes singularities as generative structures whose formation and bursts contribute to the evolving geometry of the universe.

Authors Note

The following reflection describes the moment in which the central idea of this work became clear to me

Here I am again having moments of deep thought. As I have said before this happens at the most unusual times. I say this because I was in my garden planting sweet corn, green beans, and squash when it happened. The answer to a loose end I had not conceived when I was writing The Three Realms of Existence. It was the component of the creation of a singularity within the Deep Liquid Metallic Hydrogen Realm. I recognized this as a loose end. I had it in deep thought for quite a long time and yet the formation eluded me. Until today a phrase formed within my mind.

In the deep realm the creation of a singularity is an effect of the near mass pressure within the deep realm causing gravitational convergence that forms a spherical singularity structure." I stopped took a deep breath and a moment of clarity overwhelmed me. Something I have felt many times before. What I will be describing in this work is not conventional by any means. My thinking is mechanical and structural. In this I am not just describing phenomena; I am trying to follow the physical chain of events step by step, the way an engineer studies a machine.

In my mind I keep asking:

- Where does it form?
- What pressure state exists before formation?
- What instability starts the process?
- What geometry locks the structure?
- What happens to matter, light, and information once inside?
- How does the system eventually release what it has stored?

So my mind naturally builds cycles and mechanisms, not isolated events.

That's why the pieces of this theory connect the way they do:
Deep realm pressure
→ instability node
→ near-mass threshold
→ curvature locking
→ singularity formation
→ system stabilizes
→ transition into the Relative Quantum Realm (Our Part of the Universe)

→ geometric capture and quantum stasis
→ accumulation of matter, light, and information
→ confinement failure
→ burst
→ wavefront propagation shaping cosmic structure

I am always looking for the mechanism that closes the loop.

And something else about my thinking became clearer: I use geometry as the governing principle. Instead of treating gravity, light, and matter as separate problems, I look at how curvature organizes them all at once. That's why ideas like:

- geometric capture of light
- helical ("rifled") gravity wells that impart angular momentum to create rotating singularities
- smooth gravity wells for non-rotating singularities
- spherical vs helical bursts
- curvature locking
- near stasis state for preservation of mass and energy quantum stasis feel natural to me — they're all expressions of geometry controlling motion.

I'll respond carefully here, because now I am describing a core statement inside my model, not just strategy or placement. What I've written expresses an internal mechanism in my framework: In the deep realm, near-mass pressure produces gravitational convergence, leading to formation of a singularity.

Let's Separate Three Layers so the Idea Stays Clear and Strong

- Inside my theoretical picture Within this model, the logic appears to be: a deep realm exists in a near-equilibrium state, mass-energy density approaches an extreme threshold, pressure and curvature concentrate, convergence occurs, a bounded structure — a singularity — forms. The singularity is not an instantaneous point of collapse, but an emergent structure produced by environmental conditions. Conceptually, this treats singularities as formed states, not mathematical endpoints. That is internally coherent as a narrative mechanism.
- Comparison with established physics In standard General Relativity: singularities appear when gravitational collapse drives curvature toward infinity, pressure and density increase during collapse, but the theory itself breaks down at the singularity. Physicists have already suspected the point singularity signals incomplete physics, which is why many quantum-gravity models explore alternatives such as: finite cores, Planck-scale structures, bounce models, regular (non-point) black holes. So my instinct of — replacing a zero-point with a structured formation process — sits in a region people actively investigate, though with very different mathematical tools [2].
- The key challenge moving forward For an idea like this to become scientifically engageable, readers naturally ask: What defines "near-mass pressure" quantitatively? What triggers convergence? What stabilizes the singularity boundary? What observable consequence differs from standard collapse? Those questions don't invalidate the idea — they're simply the bridge between conceptual description and physical model. A constructive way to phrase it A physics-aligned wording might be something like:

Within the deep realm, increasing energy density drives gravitational convergence, producing a bounded singular structure rather than a mathematical point. That keeps my intended meaning while framing it in terms familiar to readers trained in gravitational physics. What I am doing here is trying to describe singularity formation as a phase transition of spacetime, rather than a geometric failure. That's the conceptual heart of what I am expressing.

References

1. Nothstine, D. (2026b). The three realms of existence [Preprint].
2. Hawking, S. W., & Ellis, G. F. R. (1973). The large scale structure of space-time. Cambridge University Press.
3. Einstein, A. (1916). Die Grundlage der allgemeinen Relativitätstheorie. Annalen der Physik, 354(7), 769–822.
4. Nothstine, D. (2026a). Dark Star II [Preprint]. Zenodo.
5. Nothstine, D. (2025). Genesis: A new vision of our cosmos we live in, Parts I–IV [Preprint].
6. Misner, C. W., Thorne, K. S., & Wheeler, J. A. (1973). Gravitation. W. H. Freeman.
7. Wald, R. M. (1984). General relativity. University of Chicago Press.